

# Ultimate truth *vis à vis* stable truth

P.D. Welch

*School of Mathematics, University of Bristol, England.*

November 7, 2003

**Abstract.** We show that the set of *ultimately true* sentences in Hartry Field's Revenge-immune solution to the semantic paradoxes is recursively isomorphic to the set of *stably true* sentences obtained in Hans Herzberger's revision sequence starting from the null hypothesis. We further remark that this shows that a substantial subsystem of second order number theory is needed to establish the semantic values of sentences over the ground model of the standard natural numbers:  $\Delta_3^1\text{-CA}_0$  (second order number theory with a  $\Delta_3^1$ -Comprehension Axiom scheme) is insufficient.

**Keywords:** theories of truth, fixed points, revision sequence, constructible sets, admissibility.

## 1. Introduction.

In (Field, 2003) Field constructs a system of a theory of truth over ground models  $\mathcal{M}$ , with an additional conditional operator  $\rightarrow$  in the language. We shall here use for illustrative purposes  $\mathcal{M} = \mathbb{N}$ , the standard model of arithmetic, with  $\mathcal{L}$  the appropriate arithmetical language using the connectives  $\wedge, \vee, \neg$  and the two quantifiers  $\exists, \forall$ ;  $\mathcal{L}^+$  will be, just as for him, this language augmented with the  $\rightarrow$  connective, to be considered as an operator, and  $Tr$  as a predicate.)

The construction in (Field, 2003) Section 2 is in ordinal stages, where each successor ordinal stage performs an entire construction of the “next strong Kleene fixed point” *à la* Kripke. Thus the transition of  $\alpha \mapsto \alpha + 1$  involves the usual Kripkean monotonic construction over an assignment of values amongst  $\{0, 1, \frac{1}{2}\}$  to the “conditionals” of the form  $|A \rightarrow B|_\alpha$ . (We are suppressing the secondary ordinal subscript of this latter intermediate construction, which Field calls the “mini-stages” - and also for the most part suppresses.) The latter assignments are determined by semantic values given at strictly previous (full) stages. We repeat his Clause 8 to illustrate:

$$|A \rightarrow B|_\alpha = \begin{cases} 1 & : \text{ iff } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha)(|A|_\gamma \leq |B|_\gamma), \\ 0 & : \text{ iff } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha)(|A|_\gamma > |B|_\gamma), \\ \frac{1}{2} & : \text{ otherwise.} \end{cases}$$

Although the motivations are entirely different (and this cannot be emphasised enough perhaps) the system here has striking *structural* similarities with the revision theory of truth using Herzberger style revision sequences (*cf.* (Herzberger, 1982b), (Herzberger, 1982a)). For a revision sequence, know-

ing all truth values at some stage  $\alpha + 1$ , we can read off which truth value was assigned to  $B$  at stage  $\alpha$  from  $\text{Tr}(B)$ . In Field's system we have the same effect: the "1 or non-1" semantic value of  $B$  at the previous stage  $\alpha$  is imported for the next stage  $\alpha + 1$  via  $|\top \rightarrow B|_{\alpha+1}$ : knowing these semantic values means we know what happened before. There are similar, but differing, considerations for limit stages. For Herzberger, at limit stages the extension of  $Tr$  is determined through similar looking rules to the first two clauses in the above definition, however there is no value of  $\frac{1}{2}$ , but in this third case an (arbitrary?) assignment of 0. In a single Herzberger sequence one can look at the set of *stable sentences*, that is, those that from some point on have a set truth-value. (We emphasise that we are *not* considering *categorical truth sets* in the style of Gupta and Belnap (Gupta and Belnap, 1993), where a global averaging is obtained by considering all possible starting hypotheses for the Herzberger revision process.)

The values  $|A \rightarrow B|_{\alpha}$  are continuous at limits (*cf.* his Continuity Lemma for Conditionals). It then makes sense to define  $\|A\|$  as the ultimate value of  $|A|_{\alpha}$ :

$$\|A\| = \begin{cases} 1 & : \text{ iff } (\exists\beta)(\forall\gamma \geq \beta)(|A|_{\gamma} = 1), \\ 0 & : \text{ iff } (\exists\beta)(\forall\gamma \geq \beta)(|A|_{\gamma} = 0), \\ \frac{1}{2} & : \text{ otherwise} \end{cases}$$

Field then asks if there are *acceptable points*, that is ordinals  $\Delta$  so that for any  $A$  we have that  $\|A\| = |A|_{\Delta}$ . Section 3 of (Field, 2003) is devoted to proving that there are such. For the paper it is sufficient for him to demonstrate the existence of such points but he adds: "I believe a more informative proof should be possible, which would show among other things that acceptable fixed points ... occur before  $\Omega$ ." Here  $\Omega$  is the next initial ordinal of cardinality greater than that of the underlying model  $\mathcal{M}$  that is being discussed - so here for us this is  $\omega_1$ . In fact his proof is simple and direct, but again adds that "the price of its simplicity is that the proof is less informative than one might like about the way the values of sentences change as the level increases towards an acceptable point." The purpose of this note is to give a further proof (well, in fact 3 of them) for the existence of acceptable points (which are all below  $\Omega$ ). Probably none of them are "informative" in the latter sense he might like, and we are not sure that there could be anything very illuminating *in general*, meaning for the "general" sentence  $A$ , given the complexity of the notions involved. He shows that the least acceptable point is greater than a particular recursive ordinal  $\lambda_0$  that is used in his iterations of a "determinately true" operator  $D$ . Theorem 2 below shows that the process of calculating semantic values is *necessarily* complicated, and thus "how" a general sentence achieves its ultimate value involves looking at the theory of quasi-inductive definitions, or equivalently at the levels of the Gödel constructible hierarchy up to  $L_{\zeta}$  - the first  $\Sigma_2$ -extendible level. We show (Theorem 3) that the least acceptable

point is in fact exactly this  $\zeta$ , and so far beyond the first non-recursive ordinal  $\omega_{1ck}$ . Remark (ii) below also indicates the strength of the proposed system: any system of analysis in which we can establish these eventual semantic values is highly impredicative.

**DEFINITION 1.** (i) *The set of ultimate truths*  $D =_{df} \{\ulcorner A \urcorner : \|A\| = 1\}$ .  
(ii) *The set of (Herzberger) stable truths*  
 $H =_{df} \{\ulcorner A \urcorner : A \text{ is true in Herzberger's revision sequence starting from the null hypothesis}\}$ .

In the above by “the null hypothesis” we mean the hypothesis that all sentences are false, (or equivalently for our purposes this could be taken as “all sentences are true”: any recursive starting distribution of truth values would serve equally well). For the notion and properties of Herzberger’s revision sequence the reader may consult that author’s papers already cited, or the account in (McGee, 1991).

**THEOREM 1.** *D is recursively isomorphic to the set H.*

This set  $H$  has been noted to be equivalent to other sets of integers, defined independently. We shall comment on this further in Section 3, but briefly, if  $Q$  is the complete arithmetic quasi-inductive set, if  $\tilde{O}$  is the  $\Sigma_2$ -truth set of  $L_\zeta$  (where  $\zeta$  was named above), and if  $S$  is the set of (codes of) Turing programs that have eventually some constant value on their output tape, when allowed to run transfinitely (in the formalism of (Hamkins and Lewis, 2000)) then we have:

**Fact 1** *The following are recursively isomorphic:*

- (i)  $S$  ; (ii)  $\tilde{O}$  ; (iii)  $Q$  ; (iv)  $H$ .

Our third method of demonstrating the existence of acceptable points, is in fact directed to adding a fifth set  $D$  to this list. The following theorem is what we shall actually prove:

**THEOREM 2.** *Let D be the set of ultimate truths in Field's system. Then D is recursively isomorphic to  $\tilde{O}$ .*

The statement of the abstract, Theorem 1, then follows from the Fact quoted above. We note the following further consequences:

**Remark (i)** Using the above Fact 1 and Theorem 2 the set of ultimately true sentences over arithmetic is a *complete quasi-inductive set*; hence Field’s logic  $\models_{LCC}$  is not axiomatisable.

**Remark (ii)** The “conservativeness proof” of (Field, 2003) Sect.6 (there performed in *ZFC*) can be effected in a sufficiently strong fragment of second

order number theory, but not in  $KP$  (Kripke-Platek set theory) nor  $\Delta_3^1\text{-}CA_0$ . (For the latter theory see (Simpson, 1999).)<sup>1</sup>

Similar results will hold for countable ground models  $\mathcal{M}$  that are *acceptable* (this time in the sense of (Moschovakis, 1974)), that is, the least (Fieldian) acceptable point for  $\mathcal{M}$  will be the least  $\Sigma_2$ -extendible ordinal relative to  $\mathcal{M}$ , and hence will be less than  $|\mathcal{M}|^+$ , the least cardinal greater than that of  $\mathcal{M}$ . We shall then have (in the appropriate language, and in the appropriate sense) that  $D_{\mathcal{M}}$  is equivalent to  $H_{\mathcal{M}}$ .

The proof of Theorem (in the hard direction showing that  $\tilde{O}$  is (1-1) reducible to  $D$ ) is essentially a demonstration that Field's system carries hidden within it a construction of the Gödel hierarchy up to the level  $\zeta$ . We can show that at stage  $\alpha$  in his construction we have a uniform way of recovering certain  $\Sigma_2$ -truth sets of the levels  $L_\alpha$  of this hierarchy, whenever the latter is a model of a sufficient amount of set theory. Interestingly it seems to depend on this strength. For other levels we do not have a uniform way of doing this: although Field's clauses are uniform for all limit ordinals, the recovery of the truth sets for the  $L_\alpha$  can be obtained from  $\{A : |A|_\alpha = 1\}$ , but not always *uniformly* so.<sup>2</sup> Happily below  $\zeta$  there are many  $\alpha$  where  $L_\alpha$  models this theory, including  $\zeta$  itself; this allows us then to exploit the uniformity at  $\zeta$ , in order to find the (1-1) function decoding  $\tilde{O}$  from  $D$ .

We make a brief remark on our result concerning *recursive isomorphism* (this will be explicitly defined below). One might hope for some more perspicacious, or 'natural' translation of Herzberger stable truths directly into Field's 'ultimate truths', and hopefully *vice versa*. We think this unlikely. We do not really see one in either direction: this is to do with the fact that the process of revision according to revision sequences, has more in common with the supervaluation approach than the Kleene strong jump that Field employs in his 'mini-stages': direct translations seem to be ruled out by the incomparability of the methods. One should also remember however, that *all we ever do* with theories of truth employing a system of gödel numbering over a model such as the standard model of arithmetic, is to build a truth set using that particular coding scheme. There are infinitely many choices of (sensible) coding scheme, all of which only yield identical truth sets *up to recursive isomorphism*. However, even bearing this in mind, there seems a difficulty in finding a meaning preserving translation or mapping, between the stable truths and the ultimate truths.<sup>3</sup>

## 2. The existence of acceptable points

*First demonstration:* The construction of (Field, 2003) Section 2 is performed within  $ZFC$  using the ground model  $\mathcal{M}$ . It is in fact a very absolute construction, and absolute for transitive models of  $ZFC^-$ , that is  $ZFC$  without the

power set axiom (which is not used). Hence if  $H$  is a transitive set,  $\alpha \in H$  and  $H \models ZFC^-$  then  $H \models “|A|_\alpha = j”$  iff  $|A|_\alpha = j$ . Suppose, without loss of generality, that  $\mathcal{M}$  is countable. By the Collection Axiom we may assume that there is some ordinal  $\mu_0$  so that for all  $A \in \mathcal{L}^+$ , if  $|A|_\gamma = j$ , for all sufficiently large  $\gamma$ , then there is an  $\delta_A < \mu_0$  so that  $\forall \beta > \delta_A |A|_\beta = |A|_{\delta_A} = j$ . For the other  $A$ , ie those with  $\|A\| = \frac{1}{2}$ , which alternate value unboundedly in the ordinals, we can again appeal to Collection to find, for any ordinal  $\nu \geq \mu_0$ , some cardinal  $\mu \geq \nu$  so that if, say,  $A$  alternates its value unboundedly in the ordinals, it does so unboundedly below  $\mu$  in particular. (This already shows that the class of acceptable points is unbounded in the class of all ordinals.) But further, if  $H_\mu$  is the set of sets of hereditary cardinality less than  $\mu$ , then  $H_\mu$  is a  $ZFC^-$  model. Now by the Löwenheim-Skolem theorem let  $X \prec H$  be a countable elementary submodel of  $H$  with  $\mathcal{M} \in X$ . Let  $N$  be a transitive set isomorphic to  $X$ . (Such an  $N$  exists by the Mostowski collapsing Lemma.) Let  $\Delta < \omega_1$  be the ordinal height of  $N$ . Then  $\Delta$  is an acceptable point: one may easily verify that  $|A|_\Delta = |A|_\mu$  for any  $A$ .

By modifying the argument one may show that there is in fact a closed and unbounded set  $E \subseteq \omega_1$  of acceptable ordinals.

*Second demonstration:* One could directly show that the construction, and argument that there are acceptable points, can be performed not just in  $ZFC^-$  but in much weaker theories, for example, Kripke-Platek (KP) augmented with a  $\Sigma_2$ -Separation axiom, or even in second order number theory with  $\Pi_3^1$  Comprehension. (We aren't advocating actually *doing* this; that it could be done will follow from the third demonstration). If this is granted, then we can find countable models of such theories for which the construction is absolute. That it *cannot* be done in much weaker theories is also a concomitant of the theorem.

*Third demonstration:* This contains the heart of this note. We shall define an ordinal  $\zeta$ . To see what this is we make the following observations. The  $\Sigma_2$  nature of the defining clauses of  $\|A\|$ , together with the absolute nature of the construction, hint not only (a) that the definition of the function  $\|\cdot\|$  can be construed as a valuation coming from a quasi-inductive definition but (b) that the first repeat point, that is the first acceptable point, is  $\zeta$  - this latter ordinal has been characterised as the first  $\Sigma_2$ -extendible ordinal ((Burgess, 1986) Sect.14), that is the first ordinal  $\zeta$  so that  $L_\zeta$  has a proper  $\Sigma_2$ -elementary end extension.<sup>4</sup> Equivalently it is also the first place where all Infinite Time Turing machine (ITTM) computations have either halted or have entered an infinite loop ((Welch, 2000) Thm. 2.1). In the next theorem  $\Sigma$  is the periodicity, or ordinal length, of that loop, or, equivalently, the ordinal height of that smallest  $\Sigma_2$ -end extension of  $L_\zeta$ . We shall prove:

**THEOREM 3.** *Over the ground model  $\mathcal{M} = \mathbb{N}$  of arithmetic, the first acceptable point  $\Delta_0 = \zeta$ . The class of acceptable points above  $\zeta$  is:  $\{\Sigma.\rho \mid \rho \in On\}$ .*

To state explicitly at the outset, we shall reserve the word *recursive* to have its usual standard meaning throughout this note. The notion of  $P \leq_T Q$  “ $P$  is recursive in  $Q$ ”, for sets of integers  $P, Q$ , says simply that  $\chi_P$ , the characteristic function of  $P$ , is recursive in an oracle for  $Q$ ; alternatively put, there is an index  $e \in \omega$  so that  $\chi_P$  is the  $e$ 'th function recursive in  $Q$ :  $\chi_P = \{e\}^Q$ . To say that two sets  $P, Q$  are *recursively isomorphic* (written “ $P \equiv Q$ ”) is to say that there is a total recursive bijection  $F : \mathbb{N} \longleftrightarrow \mathbb{N}$  with  $\forall n \ n \in P \iff F(n) \in Q$ . For such  $P, Q$  membership questions about  $P$  can be converted by a pencil and paper algorithm to membership questions about  $Q$  and conversely. A weaker notion, written  $P \leq_1 Q$ , is that “ $P$  is ( $I$ - $I$ ) reducible to  $Q$ ”, that is, there is a recursive injection  $F : \mathbb{N} \longrightarrow \mathbb{N}$  with  $\forall n \ n \in P \iff F(n) \in Q$ . Here we could only paraphrase by saying that membership questions “ $n \in P$ ?” can be converted into the questions “ $F(n) \in Q$ ?” again *via* such an algorithm. Hence in this situation, as a slogan: “ $Q$  is at least as complicated as  $P$ ”. An effective version of the proof of the classical Cantor-Schröder-Bernstein theorem due to Myhill ((Rogers, 1967) Theorem (VI)) shows that  $P \equiv Q$  iff  $P \leq_1 Q$  &  $Q \leq_1 P$ .

We introduce some further definitions.

**DEFINITION 2.** *For  $x \in L$ , we let  $\rho_L(x) =_{df}$  the least  $\alpha$  such that  $x \in L_{\alpha+1} \setminus L_\alpha$ .*

For  $E$  a class of ordinals, let  $E^*$  denote  $E$  together with the set of its limit points.  $E^*$  is thus the *closure* of  $E$ .

**DEFINITION 3.** (i) *Let  $ADM = \{\alpha \mid \langle L_\alpha, \in \rangle \models KP\}$ .*

(ii) *Let  $\langle \tau_l \mid l < \omega_1 \rangle$  enumerate  $ADM^* \cap \omega_1$ .*

Here “KP” denotes Kripke-Platek set theory in the language  $\mathcal{L}_{\dot{\in}}$ . If  $\alpha \in ADM$  it is called *admissible*. Then  $\tau_0 = \omega, \tau_1 = \omega_{1ck}$ , where the latter is the first non-recursive ordinal. Note that  $ADM$  is not closed:  $\tau_\omega$  is not admissible.

We shall make use of the following facts:

**LEMMA 1.** (i)  $\zeta = \tau_\zeta$ ;  $\Sigma = \tau_\Sigma$ .

(ii)  $\forall \mu < \Sigma \neg \exists \nu \ L_\nu \prec_{\Sigma_2} L_\mu$ ,

((i) that  $\tau_\xi = \xi$  is true for any  $\Sigma_2$ -extendible ordinal  $\xi$ ; (ii) uses the leastness assumption on  $\zeta$  and so also on  $\Sigma$ .) These can be deduced from the reflection properties enjoyed by  $\Sigma_2$ -extendibles.

DEFINITION 4. Let  $C \subseteq \omega$ . We let  $A^{1,C} =_{df}$  the complete  $\Sigma_1$ -theory of  $\langle L_{\omega_{1ck}^C} [C], \in, C \rangle$ .

$A^{1,C}$  may thus be identified with a set of integers coding  $\Sigma_1$  sentences in the language  $\mathcal{L}_{\dot{\in}, \dot{C}}$ . This set has itself a  $\Sigma_1$ -definition over the structure  $\langle L_{\omega_{1ck}^C} [C], \in, C \rangle$ , which is uniform in  $C$  (that is, the defining formula is independent of the choice of set  $C$ ). Here  $\omega_{1ck}^C$  is the first ordinal not recursive in  $C$ . (We shall henceforth drop the “ck” when considering the first ordinal not recursive in some  $C$ ; we shall keep it for the first non-recursive ordinal  $\omega_{1ck}$ ; the first uncountable cardinal is thus  $\omega_1$ , to avoid any confusion.)

DEFINITION 5. For  $\iota < \omega_1$  let  $A_\iota^2 = A_\iota =_{df}$  the complete  $\Sigma_2$ -theory of  $\langle L_{\tau_\iota}, \in \rangle$ .

It is thus  $\tilde{O} =_{df} A_\zeta$  which we wish to show is recursively isomorphic to  $D$ .

PROPOSITION 4. (i) For  $\iota < \Sigma$   $\rho_L(A_\iota) = \tau_\iota$ ; (ii)  $A_\Sigma = A_\zeta$  hence  $\rho_L(A_\Sigma) = \zeta = \tau_\zeta$ ; (iii)  $\omega_1^{A_\iota} = \tau_{\iota+1}$ .

**Proof:** (i) The  $\Sigma_2$ -satisfaction relation for  $\langle L_{\tau_\iota}, \in \rangle$  is  $\Sigma_2$ -definable over this structure (cf, for example, (Devlin, 1984) Lemma VI.1.15), and thus so is  $A_\iota$ . Hence  $\rho_L(A_\iota) \leq \tau_\iota$ . Now assume that  $\iota < \Sigma$ . Then for no  $\xi$  do we have  $L_\xi \prec_{\Sigma_2} L_{\tau_\iota}$  (by Lemma 1 (ii) above). It is thus the case that every  $x \in L_{\tau_\iota}$  has a parameter free  $\Sigma_2$ -definition. In other words, the  $\Sigma_2$ -Skolem Hull of  $\emptyset$  inside  $L_{\tau_\iota}$  is all of  $L_{\tau_\iota}$  itself. We may put these facts together and show that there is a (partial) function  $F : \subseteq \omega \rightarrow L_{\tau_\iota}$  which is onto and  $\Sigma_2$ -definable over  $\langle L_{\tau_\iota}, \in \rangle$ . Now, however the diagonal set  $E$  defined by:  $n \in E \iff n \in \text{dom}(F) \wedge n \notin F(n)$  is itself  $\Sigma_2$ -definable over  $\langle L_{\tau_\iota}, \in \rangle$  and hence is in  $L_{\tau_{\iota+1}}$ . By the usual Russell style diagonal argument we cannot have that  $E \in L_{\tau_\iota}$ . Hence  $\rho_L(E) = \tau_\iota$ .  $E$  is recursive in  $A_\iota$  as the latter codes up all  $\Sigma_2$ -truths of  $\langle L_{\tau_\iota}, \in \rangle$ . Thus  $\rho_L(A_\iota) \geq \tau_\iota$  (for otherwise, we could construct  $E$  from  $A_\iota$  before stage  $\tau_\iota$  in the  $L$ -hierarchy).

For (ii): here  $\Sigma$  is the very least point where there is some  $\zeta < \Sigma$  with  $L_\zeta \prec_{\Sigma_2} L_\Sigma$ , and the above argument can not be applied. Further these two levels of the  $L$ -hierarchy have identical  $\Sigma_2$ -theories, i.e.,  $A_\Sigma = A_\zeta$ . Hence  $\rho_L(A_\Sigma) = \tau_\zeta$ . That  $\tau_\zeta = \zeta$  is Lemma 1 (i) above.

For (iii): Let  $\langle t_k^2 \mid k \in \omega \rangle$  recursively enumerate all  $\Sigma_2$ -skolem terms in the language  $\mathcal{L}_{\dot{\in}}$ . Then the following is a  $\Sigma_2$  sentence:

“ $t_k, t_l$  are both defined  $\wedge t_k, t_l \in On \wedge t_k < t_l$ .”

Since (by the comments in (i)) every ordinal below  $\tau_\iota$  is named by such a  $\Sigma_2$ -skolem term over the structure  $\langle L_{\tau_\iota}, \in \rangle$  we easily have that there is a wellordering  $w$  of  $\omega$  recursive in  $A_\iota$  (the latter coding up all  $\Sigma_2$  truths of this

structure), with the order type of  $w$  equal to  $\tau_\iota$ . Hence  $\omega_1^{A_\iota} > \tau_\iota$ . However for any  $B \subseteq \mathbb{N}$ ,  $B \in L_{\tau_{\iota+1}}$ , we have that  $\omega_1^B \leq \tau_{\iota+1}$ . In particular  $\omega_1^{A_\iota} \leq \tau_{\iota+1}$ ; by appealing to (i) we have then the desired result (iii). **QED**

The  $\Sigma_1$ -Separation scheme (see, e.g. (Barwise, 1975) Sect. I.9) is the infinite set of axioms which are the universal closure of axioms of the form:

$$\exists u \forall v (v \in u \longleftrightarrow v \in w \wedge \varphi(v))$$

where  $\varphi(v)$  is any  $\Sigma_1$ -formula (possibly with other set variables). For us, what matters is the following relatively easily proven fact (again see (Barwise, 1975), V 6.3 & 7.12):

**PROPOSITION 5.** *Suppose  $\beta > \omega$ .  $L_\beta \models \Sigma_1$ -Separation if and only if  $\{\gamma < \beta \mid L_\gamma \prec_{\Sigma_1} L_\beta\}$  is unbounded in  $\beta$ . If  $L_\beta \models \Sigma_1$ -Separation then  $\beta \in ADM \cap ADM^*$ .*

**PROPOSITION 6.** (cf.(Welch, 2000) Thm 2.1)  $L_\zeta \models \Sigma_1$ -Separation. <sup>5</sup>

We shall use the following notation. Let  $C_\alpha = (C_{\alpha,T}, C_{\alpha,F})$  where  $C_{0,T} = C_{0,F} = \emptyset$ . Then for  $\alpha > 0$  set:

$$\begin{aligned} C_{\alpha,T} &= \{\top \longrightarrow A : |\top \longrightarrow A|_\alpha = 1\}; \\ C_{\alpha,F} &= \{A \longrightarrow \perp : |A \longrightarrow \perp|_\alpha = 1\}. \end{aligned}$$

In the above we have omitted gödel corners, but we assume that  $C_\alpha \subseteq \mathbb{N}$ , after some suitable recursive coding, and pairing of  $C_{\alpha,T}$  with  $C_{\alpha,F}$ . Note that if we set, for  $\alpha \geq 0$ :

$$\begin{aligned} C_\alpha^+ &= (C_{\alpha,T}^+, C_{\alpha,F}^+) \\ C_{\alpha,T}^+ &= \{A : |A|_{\alpha,\Omega} = 1\}; \\ C_{\alpha,F}^+ &= \{A : |A|_{\alpha,\Omega} = 0\}, \end{aligned}$$

then, for example,  $C_{\alpha+1,T}$  is little different from  $C_{\alpha,T}^+$ , as the former is simply the set of members of the latter with “ $\top \longrightarrow$ ” inserted before them (and similarly for  $C_{\alpha+1,F}$  and  $C_{\alpha,F}^+$  *m.m.*). Note that if  $\xi$  is acceptable, then  $C_{\xi,T} = \{\top \longrightarrow A \mid A \in D\}$  and  $C_{\xi,F} = \{A \longrightarrow \perp \mid \neg A \in D\}$ . Hence  $C_\xi$  is (1-1) reducible to  $D$ .

**LEMMA 2.** *For  $\iota < \Sigma$  (i)  $A_\iota$  is arithmetic in  $C_\iota$ ; (ii)  $\omega_1^{C_\iota} = \tau_{\iota+1}$ ; (iii)  $\rho_L(C_\iota) = \tau_\iota$  ( $\iota > 0$ ); (iv) if  $L_{\tau_\iota} \models \Sigma_1$ -Separation, then  $A_\iota \leq_1 C_\iota$ ; moreover this latter clause is uniform in  $\iota$ .*

By “uniform in  $\iota$ ” in (iv) above, we mean that the (1-1) recursive function  $f: \mathbb{N} \rightarrow \mathbb{N}$  that effects the reduction of  $A_\iota$  to  $C_\iota$  is the same for each such  $\iota < \Sigma$ . Taking  $\iota = \zeta$  shows that  $\tilde{O} = A_\zeta \leq_1 C_\zeta \leq_1 D$ . By Proposition 5 the hypothesis of (iv) is false if  $\iota$  is a successor ordinal (and is also for many limit ordinals).

**Proof of Lemma:** We have mentioned the  $C_\alpha^+$  versions decorated with the plus sign, as this allows the following formulation.

PROPOSITION 7. (*Kripke*)  $C_\alpha^+$  is a complete  $\Pi_1^{1,C_\alpha}$  set of integers, (and hence by the above remark, so is  $C_{\alpha+1}$ ).

This was proven by Kripke in the case when  $\alpha = 0$ , for his original theory using strong Kleene. (A proof is in (Burgess, 1986).) By Field's definition, each  $C_\alpha^+$  is constructed as the Kripkean fixed point using this scheme, with the distribution of semantic values  $0, 1/2, 1$  assigned to the various sentences involving the  $\rightarrow$  operator. The proposition above then, is simply the relativisation of Kripke's result to the "starting distribution" coded into  $C_\alpha$ . Once we have  $C_\alpha^+$  is a complete  $\Pi_1^{1,C_\alpha}$  set of integers, we may apply a result that identifies such with the  $\Sigma_1$ -theory of the least admissible set over  $C_\alpha$  :

PROPOSITION 8. If  $D \subseteq \omega$ , and  $D^+$  is a complete  $\Pi_1^{1,D}$  set of integers, then  $D^+ \equiv \Sigma_1$ -Theory of  $\langle L_{\omega_1^D}[D], \in, D \rangle$ ; in fact there is a recursive bijection  $g : \mathbb{N} \longleftrightarrow \mathbb{N}$ , so that uniformly for all  $D$ ,

$$n \in D^+ \iff g(n) \in \Sigma_1 \cap \text{Sent} \wedge \langle L_{\omega_1^D}[D], \in, D \rangle \models g(n).$$

In particular:  $C_\alpha^+ \equiv \Sigma_1$ -Theory of  $\langle L_{\omega_1^{C_\alpha}}[C_\alpha], \in, C_\alpha \rangle$ .

The proposition is essentially obtained from the relativisations of the theorems of Kleene and Spector, (see, e.g., (Rogers, 1967), Theorems XLI & XLIV) that classify the  $\Pi_1^1$  predicates on integers as those  $\Sigma_1$ -definable over *HYP* (the latter the class of hyperarithmetical reals, together with the fact that the reals of  $L_{\omega_{1ck}}$  are those of *HYP*).

We now proceed to the proof of the clauses (i)-(iv) of the lemma. First observe that (ii) follows from (i) and (iii) using Proposition 4(iii): first we note that in general, if  $B$  is arithmetic in  $C$ , then  $\omega_1^B \leq \omega_1^C$ . Thus  $\omega_1^{C_\iota} \geq \tau_{\iota+1}$  (using (i) and Prop 4 (iii)). Further, using (iii), as  $\rho(C_\iota) = \tau_\iota$ ,  $\omega_1^{C_\iota} \leq \tau_{\iota+1}$ . Hence (ii) holds.

We proceed to prove (i),(iii),(iv) by induction on  $\iota < \Sigma$ . For  $\iota = 0$  they either hold vacuously or are trivial. Suppose now  $\iota = \eta + 1$ , and (i) to (iv) hold with  $\eta$  in place of  $\iota$ .  $A_\iota$  is the complete  $\Sigma_2$ -theory of  $\langle L_{\tau_\iota}, \in \rangle$ .

But note that  $\tau_{\eta+1} = \tau_\iota$  is the next admissible ordinal above  $\tau_\eta$  and hence is the height of the smallest admissible set containing  $C_\eta$  as an element:  $\tau_\iota = \omega_1^{C_\eta}$  (using for (ii) the inductive hypothesis.) As sets,  $L_{\omega_1^{C_\eta}}[C_\eta] = L_{\omega_1^{C_\eta}}$ . Using our nomenclature above

$$A^{1,C_\eta} = \Sigma_1\text{-Th}(\langle L_{\omega_1^{C_\eta}}[C_\eta], \in, C_\eta \rangle) = \Sigma_1\text{-Th}(\langle L_{\tau_\iota}, \in, C_\eta \rangle).$$

However the  $\Sigma_2$ -theory of this structure is simply obtained: it is (recursively isomorphic to) the Turing jump of  $A^{1,C_\eta}$ . Thus:

$$(A^{1,C_\eta})' \equiv \Sigma_2\text{-Th}(\langle L_{\tau_\iota}, \in, C_\eta \rangle).$$

However  $A_\iota$  is the "pure  $C$ -free" part of this theory, namely:

$$A_\iota = \Sigma_2\text{-Th}(\langle L_{\tau_\iota}, \in, C_\eta \rangle \cap \mathcal{L}_{\dot{\in}}).$$

Putting this together one gets:  $A_\iota \leq_T (A^{1, C_\eta})'$ , and the latter is arithmetic in  $A^{1, C_\eta} \equiv C_\eta^+ \equiv C_{\eta+1} = C_\iota$  (where the last sentence of Proposition 8 is used for the first (1-1) equivalence here.) Hence (i) holds for  $\iota = \eta + 1$ . We have seen that  $A^{1, C_\eta}$  is definable over  $\langle L_{\tau_\iota}, \in \rangle$  (using the parameter  $C_\eta$ ); hence  $\rho(A^{1, C_\eta}) \leq \tau_\iota$ . But we have just asserted that  $A^{1, C_\eta} \equiv C_\iota$ . Hence  $\rho(C_\iota) \leq \tau_\iota$  as well. But  $\iota < \Sigma$ , so we cannot have then  $\rho(C_\iota) < \tau_\iota$ , as otherwise we should also have  $\rho(A_\iota) < \tau_\iota$ , and this would contradict Proposition 4(i). This proves (iii) for  $\iota$ .

We now suppose that  $\iota < \Sigma$  is a limit, and the inductive hypotheses (i)-(iv) hold for  $\eta < \iota$ . As  $\iota < \Sigma$  we shall have that  $L_{\tau_\iota}$  is the  $\Sigma_2$ -skolem hull of  $\emptyset$  inside  $L_{\tau_\iota}$ ; that is,  $\forall x \in L_{\tau_\iota} \exists k L_{\tau_\iota} \models "x = t_k^2"$  where  $\langle t_k^2 \mid k \in \omega \rangle$  is the above recursive enumeration of all  $\Sigma_2$ -skolem terms of the language  $\mathcal{L}_{\dot{\in}}$ . We note that the form of such skolem functions can be expressed as :

$$(*) \quad t_k^2 = y \iff \exists u \forall v \psi_k(u, v, y)$$

for some  $\Sigma_0$   $\psi_k$ .

We divide into cases depending on whether  $L_{\tau_\iota}$  satisfies the  $\Sigma_1$ -Separation scheme or not. In the former we shall have a uniform (1-1) recursive way of recovering  $A_\iota$  from  $C_\iota$ ; in the latter case it is the presence of parameters in the argument that seemingly prevent a uniform reduction.

*Case 1  $\iota$  is a limit ordinal, but  $L_{\tau_\iota} \not\models \Sigma_1$ -Separation.*

Let  $\langle \varphi_i \mid i < \omega \rangle$  be a recursive enumeration of all  $\Sigma_1$  formulae of one free variable of the language  $\mathcal{L}_{\dot{\in}}$ . The case hypothesis assures us there is a " $\Sigma_1$ -parameter"  $p \in L_{\tau_\iota}$ , so that  $L_{\tau_\iota}$  is the  $\Sigma_1$ -skolem hull of the singleton set  $\{p\}$ <sup>6</sup>. By the comment on  $\Sigma_2$ -skolem hulls above, there is  $k$  so that  $p = t_k^2$ . Moreover, by the form at (\*) of the skolem terms  $t_n^2$ , there is  $\alpha_0$  so that for all  $\alpha < \iota$ ,  $\alpha_0 < \alpha \rightarrow L_{\tau_\alpha} \models "p = t_k^2"$ ; that is  $p$  is named in the same way in the structures  $L_{\tau_\alpha}$ .<sup>7</sup> We then have the following chain of equivalences.

$$\begin{aligned} "l \in A_{\tau_\iota}^1 &\iff_{df} L_{\tau_\iota} \models \varphi_l(p) \\ &\iff \exists \beta_0 > \alpha_0 \forall \beta > \beta_0 \quad L_{\tau_{\beta+1}} \models "L_{\tau_\beta} \models \varphi_l(t_k^2)" \\ &\iff \exists \beta_0 > \alpha_0 \forall \beta > \beta_0 \quad \langle L_{\tau_{\beta+1}}[C_\beta], \in, C_\beta \rangle \models "L_{\rho(C_\beta)} \models \varphi_l(t_k^2)" \end{aligned}$$

(using in the last equivalence the inductive hypothesis (iii)); the statement in quotation marks here is actually a  $\Sigma_1$  sentence in  $\mathcal{L}_{\dot{\in}, \dot{C}}$  about  $k, l$ ; so let us call this  $\sigma(k, l)$ . Note that the map  $(k, l) \mapsto \sigma(k, l)$  can be assumed recursive. Using the Proposition 8 the last equivalence becomes:

$$\begin{aligned} &\iff \exists \beta_0 > \alpha_0 \forall \beta > \beta_0 \quad g^{-1}(\sigma(k, l)) \in C_\beta^+ \\ &\iff g^{-1}(\sigma(k, l)) \in C_\iota. \end{aligned}$$

We thus have  $A_{\tau_\iota}^1 \leq_1 C_\iota$ . This implies that  $A_\iota$  is recursive in  $(A_{\tau_\iota}^1)' \leq_T C'_\iota$ . Hence (i) holds:  $A_\iota$  is arithmetic in  $C_\iota$ . This leaves (iii):  $\rho(C_\iota) \not\leq \tau_\iota$  (for otherwise  $\rho(C_\iota) < \tau_\alpha$  for an  $\alpha < \iota$ , hence  $A_\iota \in L_{\tau_\alpha}$ . This is a contradiction

as  $A_l^1 \notin L_{\tau_l}$ .) But  $C_l$  is  $\Sigma_2(L_{\tau_l})$ . Hence  $\rho(C_l) = \tau_l$  as required.

*Case 2*  $\iota$  is a limit ordinal, and  $L_{\tau_\iota} \models \Sigma_1$ -Separation.

Now there is no such parameter  $p$  as in *Case 1*. However now let  $\langle \varphi_l^2 \mid l \in \omega \rangle$  enumerate all  $\Sigma_2$ -sentences of  $\mathcal{L}_{\dot{\zeta}}$ .

$$\begin{aligned} "l \in A_l &\iff_{df} L_{\tau_l} \models \varphi_l^2 \\ &\iff \exists \alpha_0 \forall \beta \geq \alpha_0 L_{\tau_\beta} \models \varphi_l^2 \end{aligned}$$

We should justify this last equivalence: suppose  $\varphi_l^2$  is  $\exists u \forall v \psi_l(u, v)$ . ( $\implies$ ) If  $z$  is such that  $L_{\tau_l} \models \forall v \psi_l(u, z)$ , then we may choose  $\alpha_0 < \iota$  with  $z \in L_{\tau_{\alpha_0}}$ . Then  $\forall \beta \geq \alpha_0 L_{\tau_\beta} \models \forall v \psi_l(u, z)$ . ( $\impliedby$ ) Fix  $\alpha_0$ . By Proposition 5, find  $\alpha_0 \leq \gamma < \iota$  with  $L_{\tau_\gamma} \prec_{\Sigma_1} L_{\tau_\iota}$ . Now if  $L_{\tau_\gamma} \models \forall v \psi_l(u, z)$ , this is a  $\Pi_1$  sentence about  $z$  that persists upwards to  $L_{\tau_\iota}$ .<sup>8</sup> Now, assuming  $\alpha_0$  is such a  $\gamma$ , we continue, using similar reasoning:

$$\begin{aligned} &\iff \exists \alpha_0 \forall \beta \geq \alpha_0 L_{\tau_{\beta+1}} \models "L_{\tau_\beta} \models \varphi_l^2" \\ &\iff \exists \alpha_0 \forall \beta \geq \alpha_0 \langle L_{\tau_{\beta+1}}[C_\beta], \in, C_\beta \rangle \models "L_{\rho(C_\beta)} \models \varphi_l^2" \end{aligned}$$

(again using in the last equivalence the inductive hypothesis as before); now the sentence in quotes depends only on  $l$ , so let us call it  $\sigma(l)$ , again with  $l \mapsto \sigma(l)$  assumed recursive. For the same reasons, we have the last equivalence:

$$\begin{aligned} &\iff \exists \alpha_0 \forall \beta > \alpha_0 g^{-1}(\sigma(l)) \in C_\beta^+ \\ &\iff g^{-1}(\sigma(l)) \in C_\iota. \end{aligned}$$

Thus  $A_l \leq_1 C_l$  and the reduction here ( $g^{-1} \circ \sigma$ ) is independent of  $\iota$ .

QED (Lemma 2)

**Proof of Theorem 3.** As stated at the beginning of this “*third demonstration*”, we could consider running Field’s construction inside  $L$ , indeed inside  $L_\Sigma$ . The defining clauses of his semantic value assignments form a  $\Sigma_2$ -recursion. We note then:

(1)  $|A|_\zeta = |A|_\Sigma$  for any sentence  $A$  of  $\mathcal{L}^+$ . Hence  $C_\zeta = C_\Sigma$ .

Proof: Immediate from the fact that there is a uniform  $\Sigma_2$ -recursion in the sentence variable  $A$  that defines these values over any  $L_\lambda$ , if  $\lambda = \tau_\lambda$ , say, and then  $|A|_\lambda$  will be  $\Sigma_2$ -definable over  $L_\lambda$ . Focussing on  $L_\zeta$ , and  $L_\Sigma$  respectively, we have that  $L_\zeta \prec_{\Sigma_2} L_\Sigma$ , and the result then follows.

QED(1)

(2) If  $|A|_\zeta = 0$  (resp. 1) then  $\forall \alpha \in (\zeta, \Sigma) |A|_\alpha = 0$  (resp. 1).

Proof: Suppose this were false for some particular  $A$  with, say,  $|A|_\zeta = 0$ , but  $|A|_\alpha \neq 0$ . Suppose  $\beta_0 < \zeta$  is such that  $\forall \beta \in (\beta_0, \zeta) |A|_\beta = 0$ . By supposition  $L_\Sigma \models \exists \alpha > \beta_0 |A|_\alpha \neq 0$ . The latter is a  $\Sigma_1$  sentence about  $A$  and  $\beta_0$  and will reflect down to  $L_\zeta$  (as all  $\Sigma_2$ -sentences about objects in  $L_\zeta$  do). Hence  $L_\zeta \models \exists \alpha > \beta_0 |A|_\alpha \neq 0$ . However this contradicts our choice of  $\beta_0$ . QED(2)

However (1) implies that the distribution of semantic values according to Field’s clauses are exactly the same at stage  $\zeta$  and at stage  $\Sigma$ . Hence they will

be identical at stage  $\zeta + 1$  and  $\Sigma + 1$  etc., etc. The force of (2) is that any 0,1 semantic value assigned at stage  $\zeta$  to a sentence  $A$  will remain assigned *at all later stages*. Hence  $\zeta$  is an acceptable point.

(3)  $\zeta$  is the least acceptable point.

**Proof:** Suppose for a contradiction that  $\xi < \zeta$  was the least acceptable point. We claim there is a  $\theta < \zeta$  which is an acceptable point and further  $L_\theta \models \Sigma_1$ -Separation. This is because  $L_\Sigma \models$  “There exists  $\theta > \xi$  with  $C_\theta = C_\xi$  and  $L_\theta \models \Sigma_1$ -Separation” ( $\zeta$  is such). Hence there is such a  $\theta < \zeta$ . However then we have that  $C_\theta = C_\zeta$ . This is a contradiction since, by Lemma 2,  $A_\theta$  and  $A_\zeta$  are recursively embedded into  $C_\theta, C_\zeta$  via the same function. Since  $A_\theta \neq A_\zeta$  (because  $(L_\zeta, L_\Sigma)$  are the least pair of levels of the  $L$ -hierarchy with the same  $\Sigma_2$ -theories) we cannot have  $C_\theta = C_\zeta$ ! Contradiction! QED(3)

The sequence of sets of assignment values will then reappear with a periodicity  $\Sigma$ , as required for the theorem. QED(Theorem 3)

**Proof of Theorem 2** Lemma 2 shows that  $\tilde{O} \leq_1 D$ . However the argument at (1) above shows that there is a  $\Sigma_2$  formula  $\psi(v_0) \in \mathcal{L}_{\tilde{c}}$  so that  $A \in D$  if and only if  $\psi(A)$  is true at  $L_\zeta$  (equivalently  $L_\Sigma$ ). Hence  $D \leq_1 \tilde{O}$ . The result follows by the effective Cantor-Schröder-Bernstein Theorem mentioned above. QED(Theorem 2)

### 3. Further equivalences.

**Fact 1**<sup>9</sup> The following are recursively isomorphic:

- (i)  $S$ ; (ii)  $A_\zeta$ ; (iii)  $Q$ ; (iv)  $H$ .

We introduce some nomenclature (taken from (Hamkins and Lewis, 2000)) in order to talk about the class of *infinite time turing machine* (ITTM) computations. This will be the straightforward generalisation of that from ordinary recursion theory. We let  $\langle P_e \mid e \in \omega \rangle$  be an enumeration of all ITTM programs (which are no different from ordinary Turing programs, apart from the addition of a single new “*limit state*”,  $q_L$ , to which the machine enters at limit stages of time). We use the notation that “ $P_e(k) \downarrow l$ ” to mean the computation halts with output tape containing the string  $l \in 2^\omega$ . “ $P_e(k) \uparrow$ ” means that the computation halts, or eventually enters an infinite loop, but that 0 (the infinite string of zeros) remains on the output tape (even if the machine has not formally halted). Note there is, as expected, a third class of computations: it may be that the  $e$ 'th program on input  $k$  loops without any stable contents on its output tape, we denote this as usual : “ $P_e(k) \uparrow$ ”.  $S$  to follow is a generalisation of *Turing jump*.

DEFINITION 6.  $S =_{df} \{e : P_e(0) \uparrow\}$

A perhaps more understandable version of any of the sets we have been considering is the set  $Q$ , the *complete arithmetic quasi-inductive* set of integers. To see what  $Q$  is we follow Burgess (Burgess, 1986) (who distilled this definition from that of a Herzberger revision sequence), and consider the following.

Let  $\Gamma : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$  be any arithmetic operator (that is “ $n \in \Gamma(X)$ ” is arithmetic; we emphasise that  $\Gamma$  need be neither monotone nor progressive). We define the following iterates of  $\Gamma : \Gamma_0(X) = \Gamma(X); \Gamma_{\alpha+1}(X) = \Gamma(\Gamma_\alpha(X)); \Gamma_\lambda(X) = \liminf_{\alpha \rightarrow \lambda} \Gamma_\alpha(X) = \bigcup_{\alpha < \lambda} \bigcap_{\lambda > \beta > \alpha} \Gamma_\beta(X)$ . We say that  $Y \subseteq \omega$  is *arithmetically quasi-inductive* if for some such  $\Gamma$ ,  $Y$  is (1-1) reducible via a recursive function to  $\Gamma_{\mathcal{O}_n}(\emptyset)$ . The *complete* arithmetically quasi-inductive set  $Q$ , to which all others are similarly reducible, has then also the various recursively isomorphic characterisations stated in Fact 1 above. The list of recursively isomorphic sets displayed in the fact above all arise from definitions or procedures that can be seen to fall under the scheme of being arithmetically quasi-inductive. It can be shown that the set of stable truths of a Herzberger revision sequence, and Field’s set of ultimate truths, can be obtained *via* an arithmetic quasi-inductive definition.<sup>10</sup> Each ITTM program can be thought of as producing a set of 0/1 values on an infinite tape according to a *recursive* quasi-inductive definition. Moreover, by consideration of a universal ITTM program, one may see that the complete arithmetical quasi-inductive set is in fact recursively quasi-inductive (and hence so are  $D, H \dots$  etc.) A further example of an occurrence of this scheme appears in computer science: see (Kreutzer, 2002) where the author seeks to separate out various fixed point logics on finite structures, by considering essentially the same kind of infinite quasi-inductive process.

**COROLLARY 1.** *Let  $\langle A_n | n \in \omega \rangle$  be a recursive enumeration of all sentences of  $\mathcal{L}^+$ . Let  $B \subseteq \mathbb{N}$  be any arithmetical quasi-inductive set. Then there is a recursive (1-1) function  $f$  so that so that  $B = \{n : A_{f(n)} \in D\}$ .*

Perhaps unsurprisingly:

**COROLLARY 2.**  $\models_{LCC}$  is not axiomatisable.

The last part of the Remark (ii) - that certain theories are insufficient for the conservativeness result of Field’s Sect.6 - is only an observation that is a consequence of the fact that such theories cannot prove the existence of acceptable points, in particular of  $\zeta$ . The least  $\beta$ -model of  $\Delta^1_3\text{-CA}_0$  occurs as the set of the reals of some level  $L_\gamma$  of the constructible hierarchy, for an ordinal  $\gamma$  that is much smaller than  $\zeta$ . Hence we can find such a model containing no code for a wellordering along which we can perform an arithmetic quasi-inductive definition and reach a point in the ordering where we have

a repetition in the values produced. In other words such orderings have no “acceptable” points.

### Notes

<sup>1</sup> In brief, this is second order number theory with a set induction scheme, but with comprehension restricted to formulae expressible in a  $\Delta_3^1$  fashion.

<sup>2</sup> In a nutshell: we are working in a part of the constructible hierarchy, where each level has a  $\Sigma_1$ -truth set that is definable using, in general, a necessary infinite ordinal parameter. This parameter has a  $\Sigma_2$ -definition over  $L_\alpha$  but not a uniform one independent of  $\alpha$ . For  $L_\alpha$  which are models of  $\Sigma_1$ -Separation, there is in any case no such parameter from which all sets are  $\Sigma_1$ -definable. However for these latter  $L_\alpha$ , we do in fact have a uniform way of recovering a  $\Sigma_2$  truth set, as this is the “liminf” of the previous  $\Sigma_2$ -truth sets for the models  $L_\beta$  ( $\beta < \alpha$ ).

<sup>3</sup> One might further add that a similar situation pertains to the least Kripkean fixed points under the various schemes, such as weak- and strong-Kleene, and the supervaluation scheme: such sets are all recursively isomorphic - without there being any obvious recursive “translation” mechanism.

<sup>4</sup> This means that for some  $\xi > \zeta$  we have that  $L_\zeta \prec_{\Sigma_2} L_\xi$ , the subscript  $\Sigma_2$  indicating that  $\Sigma_2$  formulae with parameters allowed from  $L_\zeta$  have the same truth value in both structures.

<sup>5</sup> Actually the theorem cited shows that  $L_\zeta$  is a model of the  $\Delta_2$ -Comprehension, and  $\Sigma_2$ -Collection Schemes, but from these two the  $\Sigma_1$ -Separation scheme follows.

<sup>6</sup> The existence of such a parameter can be established as follows: suppose there were no such  $p$ . Then the  $\Sigma_1$ -skolem hull of  $\emptyset$  in  $L_{\tau_i}$ , is not all of  $L_{\tau_i}$ ; one can argue that this hull is transitive, and in fact is thus  $L_{\alpha_0}$  for some  $\alpha_0 < \tau_i$ . Hence  $L_{\alpha_0} \prec_{\Sigma_1} L_{\tau_i}$ . Now consider the  $\Sigma_1$ -skolem hull of  $\{\alpha_0\}$  in  $L_{\tau_i}$ ; again by supposition this is not all of  $L_{\tau_i}$ ; it is transitive and hence is some  $L_{\alpha_1} \prec_{\Sigma_1} L_{\tau_i}$ , with  $\alpha_0 < \alpha_1 < \tau_i$ . Continuing in this way we could create an infinite chain of such substructures of ordinal height  $\alpha_n < \tau_i$ . This would contradict Proposition 5. In such a situation as here there is a “standard  $\Sigma_1$ -parameter”  $p = p_{\tau_i}^1 \in L_{\tau_i}$ , so that  $L_{\tau_i}$  is the  $\Sigma_1$ -skolem hull of the singleton set  $\{p\}$ . Such a parameter is least in a particular wellordering on finite sequences of ordinals below  $\tau_i$ . It can be shown that in our situation this standard parameter is a single ordinal  $\pi < \tau_i$ , (perhaps just 0). For a fuller account of such parameters and hulls see (Devlin, 1984) IV.

<sup>7</sup> The point here is that if we suppose that  $L_{\tau_i} \models “p = t_k^2”$  where  $t_k^2 = p \iff \exists u \forall v \psi_k(u, v, p)$ , then if  $L_{\tau_i} \models \forall v \psi_k(z, v, p)$  for some particular  $z$  we may take  $\alpha_0$  sufficiently large so that  $z \in L_{\tau_{\alpha_0}}$ . Then for any  $\alpha > \alpha_0$  we shall have  $L_{\tau_\alpha} \models \forall v \psi_k(z, v, p)$  (as the latter is a  $\Pi_1$ , *i.e.* a universal statement). Thus  $p$  is named in the same way by  $t_k^2$  in all such structures. Note that  $p$  will have other, different,  $\Sigma_2$  names in each such structure, but they will not necessarily persist upwards to  $L_{\tau_i}$ .

<sup>8</sup> Note how this argument can fail if  $L_{\tau_i}$  is not a model of  $\Sigma_1$ -Separation: take  $i = \omega$ , and  $\varphi_i^2$  as “there is a largest admissible ordinal”. Then for all  $n \in \omega$ ,  $L_{\tau_n} \models \varphi_i^2$ , but this sentence is false in  $L_{\tau_\omega}$  (and moreover  $g^{-1}(\sigma(i)) \in C_\omega$ ).

<sup>9</sup> The equivalence of (i) and (ii) is (Welch, 2000) Thm 2.6; whilst working on this latter paper, we were unaware of the very concrete links with the much earlier work of Burgess in (Burgess, 1986). There he explicitly states the equivalence of (iii) and (iv) as Prop.13.1; but the equivalence of (ii) and (iii) is essentially his Theorem 14.1 there too.

<sup>10</sup> This is done directly for the Herzberger sequence with a null starting hypothesis in (Burgess, 1986).

## References

- Barwise, K.: 1975, *Admissible Sets and Structures*, Perspectives in Mathematical Logic. Springer Verlag.
- Burgess, J.: 1986, 'The truth is never simple'. *Journal for Symbolic Logic* **51**(3), 663–681.
- Devlin, K.: 1984, *Constructibility*, Perspectives in Mathematical Logic. Berlin, Heidelberg: Springer.
- Field, H.: 2003, 'A revenge-immune solution to the semantic paradoxes'. *Journal of Philosophical Logic* **32**(3), 139–177.
- Gupta, A. and N. Belnap: 1993, *The revision theory of truth*. Cambridge: M.I.T. Press.
- Hamkins, J. D. and A. Lewis: 2000, 'Infinite Time Turing Machines'. *Journal of Symbolic Logic* **65**(2), 567–604.
- Herzberger, H.: 1982a, 'Naive Semantics and the Liar Paradox'. *Journal of Philosophy* **79**, 479–497.
- Herzberger, H.: 1982b, 'Notes on naive semantics'. *Journal of Philosophical Logic* **11**, 61–102.
- Kreutzer, S.: 2002, 'Partial fixed point logic on infinite structures'. In: *Annual Conference of the European Association for Computer Science Logic (CSL)*, Vol. 2471 of *Lecture Notes in Computer Science*. Springer.
- McGee, V.: 1991, *Truth, Vagueness, and Paradox: An essay on the Logic of Truth*. Hackett.
- Moschovakis, Y.: 1974, *Elementary Induction on Abstract structures*, Vol. 77 of *Studies in Logic series*. Amsterdam: North-Holland.
- Rogers, H.: 1967, *Recursive Function Theory*, Higher Mathematics. McGraw.
- Simpson, S.: 1999, *Subsystems of second order arithmetic*, Perspectives in Mathematical Logic. Springer.
- Welch, P.: 2000, 'Eventually Infinite Time Turing Degrees: infinite time decidable reals'. *Journal for Symbolic Logic* **65**(3), 1193–1203.

