

Probability 1, Autumn 2015, Problem sheet 10

To be discussed on the week 7 Dec. . . 11 Dec.

Problems marked with “PrCl” are discussed in the large problem class on Thursday 3 Dec.
Mandatory HW’s are marked with “HW”, they are due on the week 14 Dec. . . 18 Dec, the latest.¹
Solutions will be available on Blackboard on the 19th Dec.

10.1 If X and Y are independent and identically distributed with mean μ and variance σ^2 , find $\mathbf{E}[(X - Y)^2]$.

HW 10.2 In my purse the number of one penny, two pence, five pence, ten pence, twenty pence, fifty pence, one pound and two pounds coins are i.i.d. Poisson random variables with parameter λ . Use linearity of expectations to compute the expectation and variance of the joint value of my coins.

10.3 We roll two fair dice, and let X to be the sum of the numbers shown, Y to be the number on the first die minus the number on the second die. Find $\mathbf{Cov}(X, Y)$. Are X and Y independent?

PrCl 10.4 The matching problem. Recall problem **3.5**: n gentlemen go out for dinner, and they leave their hats in the cloakroom. After the dinner (and several glasses of wine) they pick their hats completely randomly. Denote by X the number of gen’s who take their own hats. This time find $\mathbf{E}X$ and $\mathbf{Var}X$.

HW 10.5 5 students enter the elevator on the ground floor of the Maths Building, and they each choose one of the floors 1 . . . 4 independently and randomly. Find the expectation and variance of the number of stops the elevator makes. Use indicators for the floors.

10.6 We roll a fair die n times. Let X be the number of times we see $\begin{bmatrix} \cdot & \cdot \\ \cdot & \cdot \end{bmatrix}$ occurring and Y the number of times we see $\begin{bmatrix} \cdot & \cdot \\ \cdot & \cdot \end{bmatrix}$. Compute the correlation coefficient of these two variables.

10.7 A *graph* consists of *vertices* and *edges* that connect two different vertices. Consider a graph of n vertices, and assume that each of the $\binom{n}{2}$ possible edges between them are independently present with probability p and absent with probability $1 - p$. (This is called the Erdős-Rényi random graph.) The *degree* D_i of vertex i is the number of edges with one end being i ($i = 1, 2, \dots, n$).

(a) What is the distribution of D_i ?

(b) Compute the correlation coefficient of D_i and D_j for $i \neq j$. *HINT: Write D_i and D_j in terms of indicators $I_{\ell k}$ of the presence of the edge between vertices ℓ and k , where $\ell, k = 1, 2, \dots, n, \ell \neq k$.*

HW 10.8 Let X_1, X_2, \dots be i.i.d. variables with mean μ and variance σ^2 , and define $Y_n := X_n + X_{n+1} + X_{n+2}$. Calculate $\varrho(Y_n, Y_{n+j})$ for all $j \geq 0$.

10.9 Suppose $Y = aX + b$ and show

$$\varrho(X, Y) = \begin{cases} +1 & \text{if } a > 0, \\ -1 & \text{if } a < 0. \end{cases}$$

10.10 Best prediction of X based on Y .

(a) Prove Steiner’s theorem: for any $c \in \mathbb{R}$, $\mathbf{E}(X - c)^2 = \mathbf{Var}X + (c - \mathbf{E}X)^2$. (*This is the same Steiner’s Theorem as the one you might have seen in Physics about moments of inertia.*)

(b) By considering conditional expectations rather than ordinary ones, conclude that for any $c(Y)$,

$$\mathbf{E}((X - c(Y))^2 | Y) = \mathbf{Var}(X | Y) + (c(Y) - \mathbf{E}(X | Y))^2.$$

In particular, $c(Y) = \mathbf{E}(X | Y)$ makes the above display minimal.

(c) Apply \mathbf{E} on the above display (this expectation will be with respect to Y) to conclude that the choice $c(Y) = \mathbf{E}(X | Y)$ makes the square deviation $\mathbf{E}(X - c(Y))^2$ minimal among functions of Y .

10.11 The number of accidents that a person has in a given year is a Poisson random variable with parameter λ . However, suppose that the value of λ changes from person to person, being equal to 2 for 60 percent of the population, and 3 for the other 40 percent. A person is chosen at random. What is the probability that this person

(a) has no accidents this year;

(b) has exactly 3 accidents this year;

¹Details of how to hand in are to be discussed with your tutor.

(c) has exactly 3 accidents this year, given that in the previous year (s)he had none?

10.12 Z students enter the elevator on the ground floor of the Maths Building, where Z is random with $\mathbf{E}Z > 1$. They each choose one of the floors $1 \dots 4$ independently and randomly. Let X be the number of stops the elevator makes.

(a) Prove $\mathbf{E}X < \mathbf{E}Z$.

(b) Suppose now $Z \sim \text{Poi}(3)$ and find $\mathbf{E}X$.

10.13 Let X_1, X_2, \dots, X_n be i.i.d. random variables. Calculate $\mathbf{E}(X_1 | X_1 + X_2 + \dots + X_n = x)$. *HINT:* $\mathbf{E}(X_1 + X_2 + \dots + X_n | X_1 + X_2 + \dots + X_n = x)$.

PrCl **10.14** Let X be a standard normal variable, and I independent of X with $\mathbf{P}\{I = 1\} = \mathbf{P}\{I = 0\} = 1/2$. Define

$$Y := \begin{cases} X, & \text{if } I = 1, \\ -X, & \text{if } I = 0. \end{cases}$$

(a) Show that Y is also standard normal.

(b) Are I and Y independent?

(c) Are X and Y independent?

(d) Show that $\mathbf{Cov}(X, Y) = 0$.

10.15 Conditional covariance. The *conditional covariance of X and Y , conditioned on Z* is defined as

$$\mathbf{Cov}(X, Y | Z) = \mathbf{E}[(X - \mathbf{E}(X | Z)) \cdot (Y - \mathbf{E}(Y | Z)) | Z].$$

(a) Show that

$$\mathbf{Cov}(X, Y | Z) = \mathbf{E}(XY | Z) - \mathbf{E}(X | Z) \cdot \mathbf{E}(Y | Z).$$

(b) Prove the conditional covariance formula

$$\mathbf{Cov}(X, Y) = \mathbf{E}[\mathbf{Cov}(X, Y | Z)] + \mathbf{Cov}[\mathbf{E}(X | Z), \mathbf{E}(Y | Z)].$$

(c) Let $X = Y$ in this display: the conditional variance formula follows.

(d) Suppose that conditioning on Z , X and Y become independent with mean Z . Show that

$$\mathbf{Cov}(X, Y) = \mathbf{Var}Z.$$

(e) We repeatedly flip a biased coin that comes up Heads with probability p , and Tails with probability $q = 1 - p$. Denote by X and Y the length of the first and the second pure sequence, respectively. (E.g., if we flip $HHHTTH \dots$, then $X = 3, Y = 2$; or if we get $THHT \dots$, then $X = 1, Y = 2$.) Determine the following quantities: $\mathbf{E}X, \mathbf{E}Y, \mathbf{Var}X, \mathbf{Var}Y, \mathbf{Cov}(X, Y)$. *HINT: condition on the first flip.*